

C e

16.1	Sequences and Series	2
16.2	Infinite Series	13
16.3	The Binomial Series	26
16.4	Power Series	32
16.5	Maclaurin and Taylor Series	40

L a s t

In this Workbook you will learn about sequences and series. You will learn about arithmetic and geometric series and also about infinite series. You will learn how to test the for the convergence of an infinite series. You will then learn about power series, in particular you will study the binomial series. Finally you will apply your knowledge of power series to the process of finding series expansions of functions of a single variable. You will be able to find the Maclaurin and Taylor series expansions of simple functions about a point of interest.

Sequences and Series

16.1



Introduction

In this Section we develop the ground work for later Sections on infinite series and on power series. We begin with simple sequences of numbers and with finite series of numbers. We introduce the summation notation for the description of series. Finally, we consider arithmetic and geometric series and obtain expressions for the sum of n terms of both types of series.



Prerequisites

Before starting this Section you should ...

- understand and be able to use the basic rules of algebra
- be able to find limits of algebraic expressions



Learning Outcomes

On completion you should be able to ...

- check if a sequence of numbers is convergent
- use the summation notation to specify series
- recognise arithmetic and geometric series and find their sums



1. Introduction

A **sequence** is any succession of numbers. For example the sequence

$$1, 1, 2, 3, 5, 8, \dots$$

which is known as the Fibonacci sequence, is formed by adding two consecutive terms together to obtain the next term. The numbers in this sequence continually increase without bound and we say this sequence **diverges**. An example of a **convergent** sequence is the **harmonic sequence**

$$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$$

Here we see the magnitude of these numbers continually decrease and it is obvious that the sequence converges to the number zero. The related **alternating harmonic sequence**

$$1, -\frac{1}{2}, \frac{1}{3}, -\frac{1}{4}, \dots$$

is also convergent to the number zero. Whether or not a sequence is convergent is often easy to deduce by graphing the individual terms. The diagrams in Figure 1 show how the individual terms of the harmonic and alternating harmonic series behave as the number of terms increase.




Figure 1


Graph the sequence:

$$1, -1, 1, -1, \dots$$

Is this convergent?

Your solution

Answer



Not convergent.

The terms in the sequence do **not** converge to a particular value. The value **oscillates**.

A general sequence is denoted by

$$a_1, a_2, \dots, a_n, \dots$$



Now find the limit of a_n as n increases:

Your solution

Answer

$$\frac{n+2}{n(n+1)} = \frac{1 + \frac{2}{n}}{n+1} \rightarrow \frac{1}{n+1} \rightarrow 0 \quad \text{as } n \text{ increases}$$

Hence the sequence is convergent.

2. Arithmetic and geometric progressions

Consider the sequences:

$$1, 4, 7, 10, \dots \quad \text{and} \quad 3, 1, -1, -3, \dots$$

In both, any particular term is obtained from the previous term by the **addition** of a constant value (3 and -2 respectively). Each of these sequences are said to be an **arithmetic sequence** or **arithmetic progression** and has general form:

$$a, a + d, a + 2d, a + 3d, \dots, a + (n - 1)d, \dots$$

in which a, d are given numbers. In the first example above $a = 1, d = 3$ whereas, in the second example, $a = 3, d = -2$. The difference between any two successive terms of a given arithmetic sequence gives the value of d which is called the **common difference**.

Two sequences which are **not** arithmetic sequences are:

$$1, 2, 4, 8, \dots$$

$$-1, -\frac{1}{3}, -\frac{1}{9}, -\frac{1}{27}, \dots$$

In each case a particular term is obtained from the previous term by **multiplying** by a constant factor (2 and $\frac{1}{3}$ respectively). Each is an example of a **geometric sequence** or **geometric progression** with the general form:

$$a, ar, ar^2, ar^3, \dots$$

where ' a '

Find a, d for the arithmetic sequence $3, 9, 15, \dots$

Your solution

$$a = \quad d =$$

Answer

$$a = 3, \quad d = 6$$

Find a, r for the geometric sequence $8, \frac{8}{7}, \frac{8}{49}, \dots$

Your solution

$$a = \quad r =$$

Answer

$$a = 8, \quad r = \frac{1}{7}$$

Write out the first four terms of the geometric series with $a = 4, r = -2$.

Your solution

Answer

$$4, -8, 16, -32, \dots$$



3. Series

A **series** is the sum of the terms of a sequence. For example, the **harmonic series** is

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots$$

and the **alternating harmonic series** is

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

The summation notation

If we consider a general sequence

$$a_1, a_2, \dots, a_n, \dots$$

then the sum of the first k terms $a_1 + a_2 + a_3 + \dots + a_k$ is concisely denoted by $\sum_{p=1}^k a_p$.

That is,

$$a_1 + a_2 + a_3 + \dots + a_k = \sum_{p=1}^k a_p$$

When we encounter the expression $\sum_{p=1}^k a_p$ we let the index ' p ' in the term a_p take, in turn, the values $1, 2, \dots, k$ and then add all these terms together. So, for example

$$\sum_{p=1}^3 a_p = a_1 + a_2 + a_3 \qquad \sum_{p=2}^7 a_p = a_2 + a_3 + a_4 + a_5 + a_6 + a_7$$

Note that p is a **dummy** index; any letter could be used as the index. For example $\sum_{i=1}^6 a_i$, and

$\sum_{m=1}^6 a_m$ each represent the same collection of terms: $a_1 + a_2 + a_3 + a_4 + a_5 + a_6$.

In order to be able to use this 'summation notation' we need to obtain a suitable expression for the 'typical term' in the series. For example, the finite series

$$1^2 + 2^2 + \dots + k^2$$

may be written as $\sum_{p=1}^k p^2$ since the typical term is clearly p^2 in which $p = 1, 2, 3, \dots, k$ in turn.

In the same way

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots - \frac{1}{16} = \sum_{p=1}^{16} \frac{(-1)^{p+1}}{p}$$

since an expression for the typical term in this alternating harmonic series is $a_p = \frac{(-1)^{p+1}}{p}$.

Write in summation form the series



4. Summing series

The arithmetic series

Consider the finite **arithmetic series** with 14 terms

$$1 + 3 + 5 + \cdots + 23 + 25 + 27$$

A simple way of working out the value of the sum is to create a second series which is the first written in reverse order. Thus we have two series, each with the same value A :

$$A = 1 + 3 + 5 + \cdots + 23 + 25 + 27$$

and

$$A = 27 + 25 + 23 + \cdots + 5 + 3 + 1$$

Now, adding the terms of these series in pairs

$$2A = 28 + 28 + 28 + \cdots + 28 + 28 + 28 = 28 \times 14 = 392 \quad \text{so} \quad A = 196.$$

As an example

$$1 + 3 + 5 + \dots + 27 \quad \text{has} \quad a = 1, \quad d = 2, \quad n = 14$$

$$\text{So } A = 1 + 3 + \dots + 27 = \frac{14}{2}[2 + (13)2] = 196.$$

The geometric series

We can also sum a general **geometric series**.

Let

$$G = a + ar + ar^2 + \dots + ar^{n-1}$$

be a geometric series having exactly n terms. To obtain the value of G in a more convenient form we first multiply through by the common ratio r :

$$rG = ar + ar^2 + ar^3 + \dots + ar^n$$

Now, writing the two series together:

$$G = a + ar + ar^2 + \dots + ar^{n-1}$$

$$rG = ar + ar^2 + ar^3 + \dots + ar^{n-1} + ar^n$$

Subtracting the second expression from the first we see that all terms on the right-hand side cancel



Find the sum of each of the following series:

(a) $1 + 2 + 3 + 4 + \dots + 100$

(b) $\frac{1}{2} + \frac{1}{6} + \frac{1}{18} + \frac{1}{54} + \frac{1}{162} + \frac{1}{486}$

(a) In this arithmetic series state the values of a, d, n :

Your solution

$$a = \quad d = \quad n =$$

Answer

$$a = 1, \quad d = 1, \quad n = 100.$$

Now find the sum:

Your solution

$$1 + 2 + 3 + \dots + 100 =$$

Answer

$$1 + 2 + 3 + \dots + 100 = 50(2 + 99) = 50(101) = 5050.$$

(b) In this geometric series state the values of a, r, n :

Your solution

$$a = \quad r = \quad n =$$

Answer

$$a = \frac{1}{2}, \quad r = \frac{1}{3}, \quad n = 6$$

Now find the sum:

Your solution

$$\frac{1}{2} + \frac{1}{6} + \frac{1}{18} + \frac{1}{54} + \frac{1}{162} + \frac{1}{486} =$$

Answer

$$\frac{1}{2} + \frac{1}{6} + \dots + \frac{1}{486} = \frac{1}{2} \frac{1 - \frac{1}{3}^6}{1 - \frac{1}{3}} = \frac{3}{4} \left(1 - \frac{1}{3}^6\right) = 0.74897$$

Exercises



Infinite Series

1. Introduction

Many of the series considered in Section 16.1 were examples of **finite series** in that they all involved the summation of a finite number of terms. When the number of terms in the series increases without bound we refer to the sum as an **infinite series**. Of particular concern with infinite series is whether they are convergent or divergent. For example, the infinite series

$$1 + 1 + 1 + 1 + \dots$$

is clearly divergent because the sum of the first n terms increases without bound as more and more terms are taken. It is less clear as to whether the harmonic and alternating harmonic series:

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots \qquad 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

converge or diverge. Indeed you may be surprised to find that the first is divergent and the second is convergent. What we shall do in this Section is to consider some simple convergence tests for infinite series. Although we all have an intuitive idea as to the meaning of convergence of an infinite series we must be more precise in our approach. We need a definition for convergence which we can apply rigorously.

First, using an obvious extension of the notation we have used for a finite sum of terms, we denote the infinite series:

$$a_1 + a_2 + a_3 + \dots + a_p + \dots \quad \text{by the expression} \quad \sum_{p=1}^{\infty} a_p$$

where a_p is an expression for the p^{th} term in the series. So, as examples:

$$1 + 2 + 3 + \dots = \sum_{p=1}^{\infty} p \quad \text{since the } p^{\text{th}} \text{ term is } a_p = p$$

$$1^2 + 2^2 + 3^2 + \dots = \sum_{p=1}^{\infty} p^2 \quad \text{since the } p^{\text{th}} \text{ term is } a_p = p^2$$



$$\lim_{n \rightarrow \infty} S_n = S \quad (\text{say})$$

then we define the sum of the infinite series to be S :

$$S = \sum_{p=1}^{\infty} a_p$$

and we say "the series converges to S ". Another way of stating this is to say that

$$\sum_{p=1}^{\infty} a_p = \lim_{n \rightarrow \infty} \sum_{p=1}^n a_p$$



Key Point 4

Convergence of Infinite Series

An infinite series $\sum_{p=1}^{\infty} a_p$ is convergent if the sequence of partial sums

$$S_1, S_2, S_3, \dots, S_k, \dots \quad \text{in which} \quad S_k = \sum_{p=1}^k a_p \quad \text{is convergent}$$

Divergence condition for an infinite series

An almost obvious requirement that an infinite series should be convergent is that the individual terms in the series should get smaller and smaller. This leads to the following Key Point:

Which of the following series cannot be convergent?

(a) $\frac{1}{2} + \frac{2}{3} + \frac{3}{4} + \dots$

(b) $1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots$

(c) $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$

In each case, use the condition from Key Point 5:

Your solution

(a) $a_p = \lim_{p \rightarrow \infty} a_p =$

Answer

$a_p = \frac{p}{p+1} \quad \lim_{p \rightarrow \infty} \frac{p}{p+1} = 1$

Hence series is divergent.

Your solution

(b) $a_p = \lim_{p \rightarrow \infty} a_p =$

Answer

$a_p = \frac{1}{p} \quad \lim_{p \rightarrow \infty} a_p = 0$

So this series **may** be convergent. Whether it is or not requires further testing.

Your solution

(c) $a_p = \lim_{p \rightarrow \infty} a_p =$

Answer

$a_p = \frac{(-1)^{p+1}}{p} \quad \lim_{p \rightarrow \infty} a_p = 0$ so again this series **may** be convergent.

Divergence of the harmonic series

The harmonic series:

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots$$

has a general term $a_n = \frac{1}{n}$ which clearly gets smaller and smaller as n . However, surprisingly, the series is divergent. Its divergence is demonstrated by showing that the harmonic series is greater than another series which is obviously divergent. We do this by grouping the terms of the harmonic series in a particular way:

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots \quad 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \dots$$

2. General tests for convergence

The techniques we have applied to analyse the harmonic and the alternating harmonic series are 'one-off' - they cannot be applied to infinite series in general. However, there are many tests that can be used to determine the convergence properties of infinite series. Of the large number available we shall only consider two such tests in detail.

The alternating series test

An alternating series is a special type of series in which the sign changes from one term to the next. They have the form

$$a_1 - a_2 + a_3 - a_4 + \dots$$

(in which each a_i , $i = 1, 2, 3, \dots$ is a **positive** number)

Examples are:

(a) $1 - 1 + 1 - 1 + 1 \dots$

(b) $\frac{1}{3} - \frac{2}{4} + \frac{3}{5} - \frac{4}{6} + \dots$

(c) $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$

For series of this type there is a simple criterion for convergence:



Key Point 6

The Alternating Series Test

The alternating series

$$a_1 - a_2 + a_3 - a_4 + \dots$$

(in which each a_i , $i = 1, 2, 3, \dots$ are **positive** numbers) is convergent if and only if

- the terms continually decrease:

$$a_1 > a_2 > a_3 > \dots$$

- the terms decrease to zero:

$$a_p \rightarrow 0 \text{ as } p \text{ increases} \quad (\text{mathematically } \lim_{p \rightarrow \infty} a_p = 0)$$



Which of the following series are convergent?

$$(a) \sum_{p=1}^{\infty} (-1)^p \frac{(2p-1)}{(2p+1)} \quad (b) \sum_{p=1}^{\infty} \frac{(-1)^{p+1}}{p^2}$$

(a) First, write out the series:

Your solution

Answer

$$-\frac{1}{3} + \frac{3}{5} - \frac{5}{7} + \dots$$

Now examine the series for convergence:

Your solution

Answer

$$\frac{(2p-1)}{(2p+1)} = \frac{(1 - \frac{1}{2p})}{(1 + \frac{1}{2p})} \quad 1 \text{ as } p \text{ increases.}$$

Since the individual terms of the series do not converge to zero this is therefore a divergent series.

(b) Apply the procedure used in (a) to problem (b):

Your solution

Answer

This series $1 - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$ is an alternating series of the form $a_1 - a_2 + a_3 - a_4 + \dots$ in which $a_p = \frac{1}{p^2}$. The a_p sequence is a decreasing sequence since $1 > \frac{1}{2^2} > \frac{1}{3^2} > \dots$

Also $\lim_{p \rightarrow \infty} \frac{1}{p^2} = 0$

3. The ratio test

This test, which is one of the most useful and widely used convergence tests, applies only to series of **positive terms**.



Key Point 7

The Ratio Test

Let $\sum_{p=1}^{\infty} a_p$ be a series of **positive** terms such that, as p increases, the limit of $\frac{a_{p+1}}{a_p}$ equals a number L . That is $\lim_{p \rightarrow \infty} \frac{a_{p+1}}{a_p} = L$.

It can be shown that:

- if $L > 1$, then $\sum_{p=1}^{\infty} a_p$ diverges
- if $L < 1$, then $\sum_{p=1}^{\infty} a_p$ converges
- if $L = 1$, then $\sum_{p=1}^{\infty} a_p$



Example 1

Use the ratio test to examine the convergence of the series

(a) $1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots$ (b) $1 + x + x^2 + x^3 + \dots$

Solution

(a) The general term in this series is $\frac{1}{p!}$ i.e.

$$1 + \frac{1}{2!} + \frac{1}{3!} + \dots = \sum_{p=1}^{\infty} \frac{1}{p!} \quad a_p = \frac{1}{p!} \quad \therefore \quad a_{p+1} = \frac{1}{(p+1)!}$$

and the ratio

$$\frac{a_{p+1}}{a_p} = \frac{p!}{(p+1)!} = \frac{p(p-1)\dots(3)(2)(1)}{(p+1)p(p-1)\dots(3)(2)(1)} = \frac{1}{(p+1)}$$

$$\therefore \quad \lim_{p \rightarrow \infty} \frac{a_{p+1}}{a_p} = \lim_{p \rightarrow \infty} \frac{1}{(p+1)} = 0$$

Since $0 < 1$ the series is convergent. In fact, it will be easily shown, using the t!

Use the ratio test to examine the convergence of the series:

$$\frac{1}{\ln 3} + \frac{8}{(\ln 3)^2} + \frac{27}{(\ln 3)^3} + \dots$$

First, find the general term of the series:

Your solution

$$a_p =$$

Answer

$$\frac{1}{\ln 3} + \frac{8}{(\ln 3)^2} + \dots = \sum_{p=1}^{\infty} \frac{p^3}{(\ln 3)^p} \quad \text{so} \quad a_p = \frac{p^3}{(\ln 3)^p}$$

Now find a_{p+1} :

Your solution

$$a_{p+1} =$$

Answer

$$a_{p+1} = \frac{(p+1)^3}{(\ln 3)^{p+1}}$$

Finally, obtain $\lim_{p \rightarrow \infty} \frac{a_{p+1}}{a_p}$:

Your solution

$$\frac{a_{p+1}}{a_p} = \qquad \therefore \qquad \lim_{p \rightarrow \infty} \frac{a_{p+1}}{a_p} =$$

Answer

$$\frac{a_{p+1}}{a_p} = \frac{p+1}{p} \cdot \frac{1}{(\ln 3)} \cdot \frac{1}{(\ln 3)^p} \cdot (\ln 3)^p = \frac{p+1}{p} \cdot \frac{1}{(\ln 3)}$$

Now $\frac{p+1}{p} = 1 + \frac{1}{p} \rightarrow 1$ as p increases

$$\therefore \lim_{p \rightarrow \infty} \frac{a_{p+1}}{a_p} = \frac{1}{(\ln 3)} < 1$$

Hence this is a convergent series.

Note that in all of these Examples and Tasks we have decided upon the convergence or divergence of various series; we have not been able to use the tests to discover what actual number the convergent series converges to.



4. Absolute and conditional convergence

The ratio test applies to series of positive terms. Indeed this is true of many related tests for convergence. However, as we have seen, not all series are series of positive terms. To apply the ratio test such series must first be converted into series of positive terms. This is easily done. Consider two series $\sum_{p=1}^{\infty} a_p$ and $\sum_{p=1}^{\infty} |a_p|$. The latter series, obviously directly related to the first, is a series of positive terms.

Using imprecise language, it is harder for the second series to converge than it is for the first, since, in the first, some of the terms may be negative and cancel out part of the contribution from the positive terms. No such cancellations can take place in the second series since they are all positive terms. Thus it is plausible that if

Show that the series $\sum_{n=1}^{\infty} \frac{1}{n^2}$



Exercises

1. Which of the following alternating series are convergent?

$$(a) \sum_{p=1}^{\infty} \frac{(-1)^p \ln(3)}{p} \quad (b) \sum_{p=1}^{\infty} \frac{(-1)^{p+1}}{p^2 + 1} \quad (c) \sum_{p=1}^{\infty} \frac{p \sin(2p + 1)}{(p + 100)^{\frac{1}{2}}}$$

2. Use the ratio test to examine the convergence of the series:

$$(a) \sum_{p=1}^{\infty} \frac{e^4}{(2p + 1)^{p+1}} \quad (b) \sum_{p=1}^{\infty} \frac{p^3}{p!} \quad (c) \sum_{p=1}^{\infty} \frac{1}{\bar{p}}$$

$$(d) \sum_{p=1}^{\infty} \frac{1}{(0.3)^p} \quad (e) \sum_{p=1}^{\infty} \frac{(-1)^{p+1}}{3^p}$$

3. For what values of x are the following series absolutely convergent?

$$(a) \sum_{p=1}^{\infty} \frac{(-1)^p x^p}{p} \quad (b) \sum_{p=1}^{\infty} \frac{(-1)^p x^p}{p!}$$

The Binomial Series

16.3

Introduction

In this Section we examine an important example of an infinite series, the **binomial** series:

$$1 + px + \frac{p(p-1)}{2!}x^2 + \frac{p(p-1)(p-2)}{3!}x^3 + \dots$$

We show that this series is only convergent if $|x| < 1$ and that in this case the series sums to the value $(1+x)^p$. As a special case of the binomial series we consider the situation when p is a positive integer n . In this case the infinite series reduces to a **finite** series and we obtain, by replacing x with $\frac{b}{a}$, the **binomial theorem**:

$$(b+a)^n = b^n + nb^{n-1}a + \frac{n(n-1)}{2!}b^{n-2}a^2 + \dots + a^n.$$

Finally, we use the binomial series to obtain various polynomial expressions for $(1+x)^p$ when x is 'small'.



Prerequisites

Before starting this Section you should ...

- understand the factorial notation
- have knowledge of the ratio test for convergence of infinite series.
- understand the use of inequalities



Learning Outcomes

On completion you should be able to ...

-



1. The binomial series

A very important infinite series which occurs often in applications and in algebra has the form:

$$1 + px + \frac{p(p-1)}{2!}x^2 + \frac{p(p-1)(p-2)}{3!}x^3 + \dots$$



Key Point 10

The Binomial Theorem

If n is a positive integer then the expansion of $(a + b)$ raised to the power n is given by:

$$(a + b)^n = a^n + na^{n-1}b + \frac{n(n-1)}{2!}a^{n-2}b^2 + \dots + b^n$$

This is known as the **binomial** theorem.

Use the binomial theorem to obtain (a) $(1 + x)^7$ (b) $(a + b)^4$

(a) Here $n = 7$:

Your solution

$$(1 + x)^7 =$$

Answer

$$(1 + x)^7 = 1 + 7x + 21x^2 + 35x^3 + 35x^4 + 21x^5 + 7x^6 + x^7$$

(b) Here $n = 4$:

Your solution

$$(a + b)^4 =$$

Answer

$$(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4.$$

Given that x is so small that powers of x^3 and above may be ignored in comparison to lower order terms, find a quadratic approximation of $(1 - x)^{\frac{1}{2}}$ and check5(quad7comp)-1(a)20



Answer

$$(1 - x)^{\frac{1}{2}} = 1 - \frac{1}{2}x + \frac{\frac{1}{2}(-\frac{1}{2})}{2}x^2 - \frac{\frac{1}{2}(-\frac{1}{2})(-\frac{3}{2})}{6}x^3 + \dots$$

Now obtain the quadratic approximation:

Your solution

$$(1 - x)^{\frac{1}{2}} \simeq$$

Answer

$$(1 - x)^{\frac{1}{2}} \simeq 1 - \frac{1}{2}x - \frac{1}{8}x^2$$

Now check on the validity of the approximation by choosing $x = 0.1$:

Your solution

Answer

On the left-hand side we have

$$(0.9)^{\frac{1}{2}} = 0.94868 \text{ to 5 d.p.} \quad \text{obtained by calculator}$$

whereas, using the quadratic expansion:

$$(0.9)^{\frac{1}{2}} \approx 1 - \frac{1}{2}(0.1) - \frac{1}{8}(0.1)^2 = 1 - 0.05 - (0.00125) = 0.94875.$$

so the error is only 0.00007.

What we have done in this last Task is to replace (or approximate) the function $(1 - x)^{\frac{1}{2}}$ by the simpler (polynomial) function $1 - \frac{1}{2}x - \frac{1}{8}x^2$ which is reasonable provided x is very small. This approximation is well illustrated geometrically by drawing the curves $y = (1 - x)^{\frac{1}{2}}$ and $y = 1 - \frac{1}{2}x - \frac{1}{8}x^2$. The two curves coincide when x is 'small'. See Figure 2:




Figure 2

Obtain a cubic approximation of $\frac{1}{(2+x)}$. Check your approximation for accuracy using appropriate values of x .

First write the term $\frac{1}{(2+x)}$ in a form suitable for the binomial series (refer to Key Point 9):

Your solution

$$\frac{1}{(2+x)} =$$

Answer

$$\frac{1}{2+x} = \frac{1}{2} \frac{1}{1+\frac{x}{2}} = \frac{1}{2} \left(1 + \frac{x}{2}\right)^{-1}$$

Now expand using the binomial series with $p = -1$ and $\frac{x}{2}$ instead of x , to include terms up to x^3 :

Your solution

$$\frac{1}{2} \left(1 + \frac{x}{2}\right)^{-1} =$$

Answer

$$\begin{aligned} \frac{1}{2} \left(1 + \frac{x}{2}\right)^{-1} &= \frac{1}{2} \left[1 + (-1)\frac{x}{2} + \frac{(-1)(-2)}{2!} \left(\frac{x}{2}\right)^2 + \frac{(-1)(-2)(-3)}{3!} \left(\frac{x}{2}\right)^3\right] \\ &= \frac{1}{2} - \frac{x}{4} + \frac{x^2}{8} - \frac{x^3}{16} \end{aligned}$$

State the range of x for which the binomial series of $1 + \frac{x}{2}^{-1}$ is valid:

Your solution

The series is valid if

Answer

valid as long as $\frac{x}{2} < 1$ i.e. $|x| < 2$ or $-2 < x < 2$



Power Series

16.4

Introduction

In this Section we consider power series. These are examples of infinite series where each term contains a variable, x , raised to a positive integer power. We use the ratio test to obtain the **radius of convergence** R , of the power series and state the important result that the series is absolutely convergent if $|x| < R$, divergent if $|x| > R$ and may or may not be convergent if $x = \pm R$. Finally, we extend the work to apply to general power series when the variable x is replaced by $(x - x_0)$.

Prerequisites

Before starting this Section you should ...

- have knowledge of infinite series and of the ratio test
- have knowledge of inequalities and of the factorial notation.

Learning Outcomes

On completion you should be able to ...

- explain what a power series is
 - obtain the radius of convergence for a power series
 - explain what a general power series is
-






Key Point 11

Convergence of Power Series

For a power series $\sum_{p=0} b_p x^p$ with radius of convergence R then

- the series converges absolutely if $|x| < R$
- the series diverges if $|x| > R$
- the series may be convergent or divergent at $x = \pm R$



For any particular power series $\sum_{p=0} b_p x^p$ the value of R can be obtained using the ratio test. We

know, from the ratio test that $\sum_{p=0} b_p x^p$ is absolutely convergent if

$$\lim_{p \rightarrow \infty} \frac{|b_{p+1} x^{p+1}|}{|b_p x^p|} = \lim_{p \rightarrow \infty} \frac{b_{p+1}}{b_p} |x| < 1 \quad \text{implying} \quad |x| < \lim_{p \rightarrow \infty} \frac{b_p}{b_{p+1}} \quad \text{and so} \quad R = \lim_{p \rightarrow \infty} \frac{b_p}{b_{p+1}} .$$



Example 2

(a) Find the radius of convergence of the series

$$1 + \frac{x}{2} + \frac{x^2}{3} + \frac{x^3}{4} + \dots$$

(b) Investigate what happens at the end-points $x = -1$, $x = +1$ of the region of absolute convergence.

**Solution**

(a) Here $1 + \frac{x}{2} + \frac{x^2}{3} + \frac{x^3}{4} + \dots = \sum_{p=0}^{\infty} \frac{x^p}{p+1}$

so

$$b_p = \frac{1}{p+1} \quad \therefore \quad b_{p+1} = \frac{1}{p+2}$$

In this case,

$$R = \lim_p \frac{p+2}{p+1} = 1$$

so the given series is absolutely convergent if $|x| < 1$ and is divergent if $|x| > 1$.

(b) At $x = +1$ the series is $1 + \frac{1}{2} + \frac{1}{3} + \dots$ which is divergent (the harmonic series). However, at $x = -1$ the series is $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$ which is convergent (the alternating harmonic series).

Finally, therefore, the series

$$1 + \frac{x}{2} + \frac{x^2}{3} + \frac{x^3}{4} + \dots$$

is convergent if $-1 \leq x < 1$.

Find the range of values of x for which the following power series converges:

$$1 + \frac{x}{3} + \frac{x^2}{3^2} + \frac{x^3}{3^3} + \dots$$

First find the coefficient of x^p :

Your solution

$$b_p =$$

Answer

$$b_p = \frac{1}{3^p}$$

Now find R , the radius of convergence:

Your solution

$$R = \lim_p \frac{b_p}{b_{p+1}} =$$

Answer

$$R = \lim_p \frac{b_p}{b_{p+1}} = \lim_p \frac{3^{p+1}}{3^p} = \lim_p (3) = 3.$$

When $x = \pm 3$ the series is clearly divergent. Hence the series is convergent only if $-3 < x < 3$.

3. Properties of power series

Let P_1 and P_2 represent two power series with radii of convergence R_1 and R_2 respectively. We can combine P_1 and P_2 together by addition and multiplication. We find the following properties:



Key Point 12

If P_1 and P_2 are power series with respective radii of convergence R_1 and R_2 then the sum $(P_1 + P_2)$ and the product $(P_1 P_2)$ are each power series with the radius of convergence being the **smaller** of R_1 and R_2 .

Power series can also be differentiated and integrated on a term by term basis:



Key Point 13

If P_1 is a power series with radius of convergence R_1 then

d



Using the known result that

$$\frac{1}{1+x} = 1 - x + x^2 - x^3 + \dots \quad |x| < 1$$

4. General power series

A general power series has the form

$$b_0 + b_1(x - x_0) + b_2(x - x_0)^2 + \dots = \sum_{p=0} b_p(x - x_0)^p$$

Exactly the same considerations apply to this general power series as apply to the 'special' series

$\sum_{p=0} b_p x^p$ except that the variable x is replaced by $(x - x_0)$. The radius of convergence of the general series is obtained in the same way:

$$R = \lim_{p \rightarrow \infty} \frac{b_p}{b_{p+1}}$$

and the interval of convergence is now shifted to have centre at $x = x_0$ (see Figure 4 below). The series is absolutely convergent if $|x - x_0| < R$, diverges if $|x - x_0| > R$ and may or may not converge if $|x - x_0| = R$.

Figure 4

Find the radius of convergence of the general power series

$$1 - (x - 1) + (x - 1)^2 - (x - 1)^3 + \dots$$

First find an expression for the general term:

Your solution

$$1 - (x - 1) + (x - 1)^2 - (x - 1)^3 + \dots = \sum_{p=0} (-1)^p (x - 1)^p$$

Answer

$$\sum_{p=0} (x - 1)^p (-1)^p \quad \text{so} \quad b_p = (-1)^p$$

Now obtain the radius of convergence:

Your solution

$$\lim_{p \rightarrow \infty} \frac{b_p}{b_{p+1}}$$



Finally, decide on the convergence at $|x - 1| = 1$ (i.e. at $x - 1 = -1$ and $x - 1 = 1$ i.e. $x = 0$ and $x = 2$):

Maclaurin and Taylor Series

16.5



Introduction

In this Section we examine how functions may be expressed in terms of power series. This is an extremely useful way of expressing a function since (as we shall see) we can then replace 'complicated' functions in terms of 'simple' polynomials. The only requirement (of any significance) is that the 'complicated' function should be *smooth*; this means that at a point of interest, it must be possible to differentiate the function as often as we please.



Prerequisites

Before starting this Section you should ...

- have knowledge of power series and of the ratio test
- be able to differentiate simple functions
- be familiar with the rules for combining power series



Learning Outcomes

On completion you should be able to ...

- find the Maclaurin and Taylor series expansions of given functions
- find Maclaurin expansions of functions by combining known power series together
- find Maclaurin expansions by using differentiation and integration



1. Maclaurin and Taylor series

As we shall see, many functions can be represented by power series. In fact we have already seen in earlier Sections examples of such a representation:

$$\frac{1}{1-x} = 1 + x + x^2 + \dots \quad |x| < 1$$
$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

2. The Maclaurin series

Consider a function $f(x)$ which can be differentiated at $x = 0$ as often as we please. For example $e^x, \cos x, \sin x$ would fit into this category but $|x|$ would not.

Let us assume that $f(x)$ can be represented by a power series in x :

$$f(x) = b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + \dots = \sum_{p=0}^{\infty} b_p x^p$$

where b_0, b_1, b_2, \dots are constants to be determined.

If we substitute $x = 0$ then, clearly $f(0) = b_0$

The other constants can be determined by further differentiating and, on each differentiation, substituting $x = 0$. For example, differentiating once:

$$f'(x) = 0 + b_1 + 2b_2x + 3b_3x^2 + 4b_4x^3 + \dots$$

so, putting $x = 0$, we have $f'(0) = b_1$.

Continuing to differentiate:

$$f''(x) = 0 + 2b_2 + 3(2)b_3x + 4(3)b_4x^2 + \dots$$

so

$$f''(0) = 2b_2 \quad \text{or} \quad b_2 = \frac{1}{2}$$



Example 4

Find the Maclaurin expansion of $\cos x$.

Solution

Here $f(x) = \cos x$ and, differentiating a number of times:

$$f(x) = \cos x, \quad f'(x) = -\sin x, \quad f''(x) = -\cos x, \quad f'''(x) = \sin x \quad \text{etc.}$$

Evaluating each of these at $x = 0$:

$$f(0) = 1, \quad f'(0) = 0, \quad f''(0) = -1, \quad f'''(0) = 0 \quad \text{etc.}$$

Substituting into $f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0) + \frac{x^3}{3!} f'''(0) + \dots$, gives:

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

The reader should confirm (by finding the radius of convergence) that this series is convergent for **all** values of x . The geometrical approximation to $\cos x$ by the first few terms of its Maclaurin series are shown in Figure 6.




Figure 6: Linear, quadratic and cubic approximations to $\cos x$

Find the Maclaurin expansion of $\ln(1+x)$.

(Note that we **cannot** find a Maclaurin expansion of the function $\ln x$ since $\ln x$ does not exist at $x = 0$ and so cannot be differentiated at $x = 0$.)

Find the first four derivatives of $f(x) = \ln(1+x)$:

Answer

$$f'(x) = \frac{1}{1+x}, \quad f''(x) = \frac{-1}{(1+x)^2}, \quad f'''(x) = \frac{2}{(1+x)^3},$$

generally: $f^{(n)}(x) = \frac{(-1)^{n+1}(n-1)!}{(1+x)^n}$

Now obtain $f(0)$, $f'(0)$, $f''(0)$, $f'''(0)$:

Your solution

$$f(0) = \quad f'(0) = \quad f''(0) = \quad f'''(0) =$$

Answer

$$f(0) = 0 \quad f'(0) = 1, \quad f''(0) = -1, \quad f'''(0) = 2,$$

generally: $f^{(n)}(0) = (-1)^{n+1}(n-1)!$

Hence, obtain the Maclaurin expansion of $\ln(1+x)$:

Your solution

$$\ln(1+x) =$$

Answer

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$



Note that when $x = 1$ $\ln 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} \dots$ so the alternating harmonic series converges to $\ln 2 \approx 0.693$, as stated in Section 16.2, page 17.

The Maclaurin expansion of a product of two functions: $f(x)g(x)$ is obtained by multiplying together the Maclaurin expansions of $f(x)$ and of $g(x)$ and collecting like terms together. The product series will have a radius of convergence equal to the **smaller** of the two separate radii of convergence.



Example 5

Find the Maclaurin expansion of $e^x \ln(1 + x)$.

Solution

Here, instead of finding the derivatives of $f(x) = e^x \ln(1 + x)$, we can more simply multiply together the Maclaurin expansions for e^x and $\ln(1 + x)$ which we already know:

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \quad \text{all } x$$

and

$$\ln(1 + x) = x - \frac{x^2}{2} + \frac{x^3}{3} + \dots \quad -1 < x < 1$$

The resulting power series will only be convergent if $-1 < x < 1$. Multiplying:

$$\begin{aligned} e^x \ln(1 + x) &= \left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \right) \left(x - \frac{x^2}{2} + \frac{x^3}{3} + \dots \right) \\ &= x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \\ &\quad + x^2 - \frac{x^3}{2} + \frac{x^4}{3} + \dots \\ &\quad + \frac{x^3}{2} - \frac{x^4}{4} \dots \\ &\quad + \frac{x^4}{6} \dots \\ &= x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{3x^4}{40} + \dots \quad -1 < x < 1 \end{aligned}$$

(You must take care not to miss relevant terms when carrying through the multiplication.)

Find the Maclaurin expansion of $\cos^2 x$ up to powers of x^4 . Hence write down the expansion of $\sin^2 x$ to powers of x^6 .

First, write down the expansion of $\cos x$:

Your solution

$$\cos x =$$

Answer

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + \dots$$

Now, by multiplication, find the expansion of $\cos^2 x$:

Your solution

$$\cos^2 x =$$

Answer

$$\begin{aligned}\cos^2 x &= \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} \dots\right) \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} \dots\right) \\ &= \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} \dots\right) + \left(-\frac{x^2}{2!} + \frac{x^4}{4!} \dots\right) + \left(\frac{x^4}{4!} \dots\right) + \dots = 1 - x^2 + \frac{x^4}{3} - \frac{2x^6}{45} \dots\end{aligned}$$

Now obtain the expansion of $\sin^2 x$ using a suitable trigonometric identity:

Your solution

$$\sin^2 x =$$

Answer

$$\sin^2 x = 1 -$$



Example 6

Find the Maclaurin expansion of $\tanh x$ up to powers of x^5 .

Solution

The first two derivatives of $f(x) = \tanh x$ are

$$f'(x) = \operatorname{sech}^2 x \quad f''(x) = -2\operatorname{sech}^2 x \tanh x \quad f'''(x) = 4\operatorname{sech}^2 x \tanh^2 x - 2\operatorname{sech}^4 x \quad \dots$$

giving $f(0) = 0, \quad f'(0) = 1, \quad f''(0) = 0, \quad f'''(0) = -2 \quad \dots$

This leads directly to the Maclaurin expansion as $\tanh x = x - \frac{1}{3}x^3 + \frac{2}{15}x^5 - \dots$



Example 7

The relationship between the wavelength, L , the wave period, T , and the water depth, d , for a surface wave in water is given by:

3. Differentiation of Maclaurin series

We have already noted that, by the binomial series,

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots \quad |x| < 1$$

Thus, with x replaced by $-x$

$$\frac{1}{1+x} = 1 - x + x^2 - x^3 + \dots \quad |x| < 1$$

We have previously obtained the Maclaurin expansion of $\ln(1+x)$:

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \quad -1 < x < 1$$

Now, we differentiate both sides with respect to x :

$$\frac{1}{1+x} = 1 - x + x^2 - x^3 + \dots$$

This result matches that found from the binomial series and demonstrates that the Maclaurin expansion of a function $f(x)$ may be differentiated term by term to give a series which will be the Maclaurin expansion of $\frac{df}{dx}$.

As we noted in Section 16.4 the derived series will have the **same** radius of convergence as the original series.

Find the Maclaurin expansion of $(1-x)^{-3}$ and state its radius of convergence.

First write down the expansion of $(1-x)^{-1}$:

Your solution

$$\frac{1}{1-x}$$

Answer

$$\frac{1}{1-x} = 1 + x + x^2 + \dots \quad |x| < 1$$

Now, by differentiation, obtain the expansion of $\frac{1}{(1-x)^2}$:

Your solution

$$\frac{1}{(1-x)^2} = \frac{d}{dx} \frac{1}{1-x} =$$

Answer

$$\frac{1}{(1-x)^2} = \frac{d}{dx}$$



Differentiate again to obtain the expansion of $(1 - x)^{-3}$:

Your solution

$$\frac{1}{(1-x)^3} = \frac{1}{2} \frac{d}{dx} \frac{1}{(1-x)^2} = \frac{1}{2} [\quad]$$

Answer

$$\frac{1}{(1-x)^3} = \frac{1}{2} \frac{d}{dx} \frac{1}{(1-x)^2} = \frac{1}{2} [2 + 6x + 12x^2 + 20x^3 + \dots] = 1 + 3x + 6x^2 + 10x^3 + \dots$$

Finally state its radius of convergence:

Your solution

Answer

The final series: $1 + 3x + 6x^2 + 10x^3 + \dots$ has radius of convergence $R = 1$ since the original series has this radius of convergence. This can also be found directly using the formula $R = \lim_{n \rightarrow \infty} \frac{b_n}{b_{n+1}}$ and using the fact that the coefficient of the n^{th} term is $b_n = \frac{1}{2}n(n+1)$.

4. The Taylor series

The **Taylor series** is a generalisation of the Maclaurin series being a power series developed in powers of $(x - x_0)$ rather



Exercises

1. Show that the series obtained in the last Task is convergent if $|x - 2| < 1$.
- 2.